

The splitting of properties of geometric images associated with affine vector field

P. Tadeyev¹, O. Kravchuk²

¹The National University of Water and Environmental Engineering, ²Khmelnitsky National University
Corresponding author. E-mail: ptadeyev@gmail.com, kravchukoa@mail.ru

Paper received 09.12.16; Accepted for publication 15.12.16.

Abstract. The authors have been constructed the splitting of the basic geometric images vector field (points, straights, hyperplanes and hyperquadrics) in transition from n-dimensional affine space to the space of affine connection. All investigations have been fulfilled in the moving coordinate system of zero order.

Keywords: vector field, splitting of geometric image, affine space, hyperplane, hyperquadric.

Introduction. Problem formation. The research of manifolds of homogenous and generalized spaces is connected with definition of invariant geometric images of affine connection (straights, points, k-dimensional planes, hyperquadrics, etc). It is also related to the research of vector fields in n-dimensional space of affine connection.

Publication review on the subject. The investigation of vector fields and associated with them chapters in equi-affine space is connected with the activities of a number of scientists. We should mention Yurii A. Aminov [1], D.A. Sintsov [2], S.S.Bushgues [3] and many others. In case of three-dimensional affine space this problem was investigated by V.V.Sluchayev [4] and Ch. Gheorgiev [5]. The bases of differential geometry of vector field in n-dimensional affine space have been built in [6], in this case accompanying moving coordinate system is chosen by means of the case when one of the vectors coincides with vector of vector field.

Besides the previous information we remark that the last mentioned work deals with building of invariant linear models: points, straights, hyperplanes and hyperquadrics.

Formation of the aims of research: to build invariant models for vector field in n-dimensional affine space with help of G.F.Laptiev [7] method.

The objective of the article is to receive the splitting of basic geometric images, vector fields in transition from n-dimensional affine space to n-dimensional space of affine connection.

Materials and methods. Classical space of affine connection $A_{n,n}$ is determined by system forms ω^α and ω_β^α satisfy the structure equations:

$$(1.1) \quad \begin{aligned} D\omega^\alpha &= [\omega^\beta \omega_\beta^\alpha] + R_{\beta\gamma}^\alpha [\omega^\beta \omega^\gamma] \\ D\omega_\beta^\alpha &= [\omega_\beta^\gamma \omega_\gamma^\alpha] + R_{\beta\gamma\delta}^\alpha [\omega^\gamma \omega^\delta], \quad (\alpha, \beta, \gamma, \delta = \overline{1, n}) \end{aligned}$$

In equations (1.1) values $R_{\beta\gamma}^\alpha$ are skewsymmetric on subscript indexes and in a set they form torsion tensor of $A_{n,n}$ space, values $R_{\beta\gamma\delta}^\alpha$ are skewsymmetric on indexes γ, δ and they form tensor of curvature of space $A_{n,n}$

The space of affine connection A_n is the fibering space which base is considered to be any differential manifold M_n , and its layers are some central affine spaces $A_n(u)$ related to affine moving coordinate system $\{A_0(u), \bar{e}_i(u)\}$.

In this case n independent first integrals u^1, u^2, \dots, u^n of fully integrated system of forms ω^i are local coordinates of point $A(u^i)$ of base M_n space $A_{n,n}$.

The forms $\omega^\alpha, \omega_\beta^\alpha$ invariantly denote infinitively close affine reflection of neighbouring local space (layer) into the given one with the help of moving coordinate system reflection

$$(1.2) \quad \begin{aligned} A(u + du) &\rightarrow dA(u) = \omega^\alpha \bar{e}_\alpha(u), \\ e_\alpha(u + du) &= e_\alpha(u) + de_\alpha(u) = \bar{e}_\alpha(u) + \omega_\alpha^\beta \bar{e}_\beta(u) \end{aligned}$$

Definition. Vector field in space $A_{n,n}$ is called the function in which every point $A(u)$ base of space $A_{n,n}$ corresponds definite vector $\vec{v}(u)$ which belongs to n-dimensional affine space $A_n(u)$ related to moving coordinate system $T_n = \{\bar{A}(u), \bar{e}_\alpha(u)\}$. This space, as it is obvious, is a layer over point $A(u)$.

System of differential equations of vector field in moving coordinate system of zero order (starting point $A(u)$ of vector field coincides with the end of vector \bar{A} , and vector \vec{v} coincides with ℓ_n have the form

$$(1.3) \quad \omega_n^\alpha = A_{n\beta}^\alpha \omega^\beta \quad (\alpha, \beta, \gamma, \dots = \overline{1, n}).$$

Continuing the system of differential equations (1.3) we'll receive the system of differential equations of fundamental object of the first order of vector field of space $A_{n,n}$ in the form

$$(1.4) \quad d\Lambda_{n\beta}^\alpha = \Lambda_{n\gamma}^\alpha \omega_\beta^\gamma - \Lambda_{n\beta}^\gamma \omega_\gamma^\alpha + \Lambda_{n\beta\gamma}^\alpha \omega^\gamma$$

where

$$(1.5) \quad \Lambda_{n[\beta\gamma]}^\alpha + \Lambda_{n\delta}^\alpha R_{\gamma\beta}^\delta + R_{n\beta\gamma}^\alpha = 0.$$

Continuing the system of differential equations (1.3) we'll have

$$(1.6) \quad d\Lambda_{n\beta\gamma}^\alpha = \Lambda_{n\delta}^\alpha \omega_\beta^\delta + \Lambda_{n\beta\delta}^\alpha \omega_\gamma^\delta - \Lambda_{n\beta\gamma}^\delta \omega_\delta^\alpha + \Lambda_{n\beta\gamma\delta}^\alpha \omega^\delta.$$

Succession of fundamental objects $\{\Lambda_{n\beta}^\alpha, \Lambda_{n\beta\gamma}^\alpha, \Lambda_{n\beta\gamma\delta}^\alpha, \dots\}$ lies on the basis of differential geometry of vector field in space $A_{n,n}$.

Remark. Apart from n-dimensional affine space tensors $\Lambda_{n\beta\gamma}^\alpha, \Lambda_{n\beta\gamma\delta}^\alpha, \dots$ loose symmetric properties on two down subscript indexes.

Results and discussion. Find differential equations of some invariant geometric objects joint to vector field.

Field of points. Consider point $P(x^\alpha)$ in affine space A_n . If \vec{P} – is radius-vector of this point, in this case related to affine moving coordinate system $(\vec{A}, \vec{e}_\alpha)$ it can be expressed with the following correspondence

$$(2.1) \quad \vec{P} = \vec{A} + x^\alpha \vec{e}_\alpha$$

After differentiation of (2.1) taking into account equation of structure, receive

$$(2.2) \quad dx^\alpha + x^\beta \omega_\beta^\alpha = x_\beta^\alpha \omega^\beta,$$

or in fixation of main parameters

$$(2.3) \quad \delta x^\alpha + x^\beta \pi_\beta^\alpha = 0.$$

Consider values

$$(2.4) \quad N^\alpha = \Lambda_{mn}^\alpha.$$

If differential equations values (2.4) have the form $dN^\alpha + N^\beta \omega_\beta^\alpha = N_\beta^\alpha \omega^\beta$ according to (2.2) they define invariant point.

Field of straights. Straight, which crosses the point A with directed vector $\vec{R} = v^\alpha \vec{e}_\alpha$ define as $l = [A, \vec{R}]$.

Conditions of invariance of straight will be

$$(2.5) \quad \delta \vec{R} = Q \vec{R}, \quad dQ = 0.$$

From the previous data

$$(2.6) \quad \delta v^\alpha + v^\beta \pi_\beta^\alpha = Q v^\alpha,$$

or

$$(2.7) \quad \begin{aligned} \delta v^i + v^j \pi_j^i &= Q v^i, \\ \delta v^n + v^i \pi_i^n &= Q v^n, \quad (i = 1, \overline{n-1}). \end{aligned}$$

Sometimes it is convenient to promote norm of vector \vec{R} in which $v^n = 1$.

Then

$$(2.8) \quad Q = v^i \omega_i^n.$$

Putting (2.8) in the first equation (2.7) we have

$$(2.9) \quad \delta v^i + v^j \pi_j^i - v^i v^k \pi_k^n = 0.$$

Thus, differential equations of invariance of straight have the form

$$(2.10) \quad dv^i + v^j \omega_j^i - v^i v^k \omega_k^n = v_\alpha^i \omega^\alpha.$$

Build values $V_{n\alpha}^\beta$ in condition of $(def // \alpha_\beta^\alpha // \neq 0)$

$$(2.11) \quad \Lambda_{n\alpha}^\gamma V_{n\gamma}^\beta = \delta_\alpha^\beta,$$

and with their help values

$$(2.12) \quad \begin{aligned} M^\alpha &= -V_{n\beta}^\alpha \Lambda_{nm}^\beta, \\ dM^\alpha &= -M^\beta \omega_\beta^\alpha + M_\gamma^\alpha \omega^\gamma. \end{aligned}$$

If differential equations of (2.12) we have the equations' structure (2.6), the pair $[A, \vec{M}]$, where $\vec{M} = M^\alpha \vec{e}_\alpha$ defines invariant straight.

Theorem 1 In differential neighborhood of the first order exists straight invariant with vector field which is determined by tensor M^α .

Field of hyperplanes. Equation of condition of invariance of hyperplane $v_\alpha x^\alpha + v = 0$ related to moving coordinate system $(\vec{A}, \vec{e}_\alpha)$ has the following form

$$(2.13) \quad \begin{aligned} \delta v_\alpha + v_\beta \pi_\alpha^\beta &= Q v_\alpha, \\ \delta v &= Q v. \end{aligned}$$

Q – a linear form, which means $dQ = 0$. Two cases are possible.

1. Hyperplane doesn't cross point A.

It's possible to put in this case $v = 1$, then $Q = 0$

Conditions of invariance of hyperplane take the form

$$(2.14) \quad \delta v_\alpha + v_\beta \pi_\alpha^\beta = 0.$$

2. Hyperplane crosses point A.

In this case $v = 0$. Conditions of its invariance have the form

$$(2.15) \quad \delta v_\alpha - v_\beta \pi_\alpha^\beta = Q v_\alpha.$$

Putting $v_n = 1$ conditions (2.15) have the form

$$(2.16) \quad \delta v_i - v_j \pi_i^j - \pi_i^n = 0.$$

Build values $g^{\alpha\beta} = N^\alpha M^\beta$.

In condition of $def // g^{\alpha\beta} // \neq 0$ introduce values $g^{\alpha\delta} g_{\gamma\beta} = \delta_\beta^\alpha$ and with their help

$$(2.17) \quad \begin{aligned} g_\alpha &= g_{\alpha\beta} N^\alpha, \\ \delta g_\alpha - g_\beta \pi_\alpha^\beta &= 0. \end{aligned}$$

If differential equations of value g_α have structure of differential equations (2.14) then these values define hyperplane which doesn't cross point A in the form $g_\alpha x^\alpha = 0$.

Theorem 2 In differential neighborhood of the first order exists invariant hiperplane of vector field which doesn't cross its beginning and which is determined by tensor g_α .

We build invariant hyperplanes with help of fundamental tensor of the second order Λ_{nji}^α . With this aim we build tensor

$$(2.18) \quad \begin{aligned} \Lambda_{n\alpha}^1 &= \Lambda_{n\alpha\beta}^\beta, \\ \delta \Lambda_{n\alpha}^1 - \Lambda_{n\beta}^1 \pi_\alpha^\beta &= 0 \end{aligned}$$

and also tensor

$$(2.19) \quad \begin{aligned} \Lambda_{n\alpha}^2 &= \Lambda_{n\beta\alpha}^\beta, \\ \delta \Lambda_{n\alpha}^2 - \Lambda_{n\beta}^2 \pi_\alpha^\beta &= 0 \end{aligned}$$

Obviously, tensors (2.18) and (2.19) for space $A_{n,n}$ are different, and in case of affine space A_n coincide. They denote invariant hyperplanes that do not cross the beginning of forming element and they are also defined by the equations:

$$(2.20) \quad \Lambda_{n\alpha}^1 x^\alpha + 1 = 0$$

$$(2.21) \quad \Lambda_{n\alpha}^2 x^\alpha + 1 = 0$$

Hyperplanes (2.20) and (2.21) are called the main hyperplanes of vector field.

Theorem 3. In case of transition from vector field of n-dimensional space A_n to vector field in n-dimensional space of affine connection the splitting of main hyperplanes takes place.

Differential conditions of invariance of hyperquadric joint to vector field.

The equation of hyperquadric relatively to any local moving coordinate system $(\vec{A}, \vec{e}_\alpha)$ has the form

$$(3.1) \quad A_{\alpha\beta} x^\alpha x^\beta + 2A_\alpha x^\alpha + A = 0.$$

Where $A_{\alpha\beta} = A_{\beta\alpha}$.

Find differential equations which can satisfy coefficients $A_{\alpha\beta}, A_\alpha$ and A quadric is invariantly connected with vector field.

If moving coordinate system $(\vec{A}, \vec{e}_\alpha)$ changes relatively to second parameters then new coordinate system equation obtains the form

$$(3.2) \quad (A_{\alpha\beta} + \delta A_{\alpha\beta})(x^\alpha - x^\gamma \pi_\gamma^\alpha)(x^\beta - x^\gamma \pi_\gamma^\beta) + (A_\alpha + \delta A_\alpha)(x^\alpha - x^\gamma \pi_\gamma^\alpha) + A + \delta A = 0$$

Conditions of invariance of hyperquadric give the opportunity to receive differential equations on the coefficients of hyperquadric:

$$(3.3) \quad \begin{aligned} \delta A_{\alpha\beta} - A_{\alpha\gamma} \pi_\beta^\gamma - A_{\beta\gamma} \pi_\alpha^\gamma &= \Theta A_{\alpha\beta}, \\ \delta A_\alpha - A_\gamma \pi_\alpha^\gamma &= \Theta A_\alpha, \\ \delta A &= \Theta A. \end{aligned}$$

Where $\Theta, -$ is linear form, where $d\Theta = 0$.

Hyperquadric doesn't cross the beginning of definite vector element.

In this case it is possible to take $A = 1$. Thus, $\Theta = 0$. Equations of invariance of hyperquadric obtain the form

$$(3.4) \quad \begin{aligned} \delta \Lambda_{\alpha\beta} - \Lambda_{\alpha\gamma} \pi_\beta^\gamma - \Lambda_{\beta\gamma} \pi_\alpha^\gamma &= 0, \\ \delta A_\alpha - A_\gamma \pi_\alpha^\gamma &= 0, \end{aligned}$$

describing them in details receive:

$$(3.5) \quad \begin{aligned} \delta A_{ij} - A_{ik} \pi_j^k - A_{kj} \pi_i^k - A_{in} \pi_j^n - A_{nj} \pi_i^n &= 0, \\ \delta A_{in} - A_{jn} \pi_i^j - A_{nn} \pi_i^n &= 0, \\ \delta A_{nn} &= 0, \\ \delta A_i - A_j \pi_i^j - A_n \pi_i^n &= 0, \\ \delta A_n &= 0, \end{aligned}$$

The hyperquadric proper can be introduced in this case by the equation

$$(3.6) \quad A_{\alpha\beta} x^\alpha x^\beta - 2A_\alpha x^\alpha + 1 = 0.$$

or wider by the equation form

$$(3.7) \quad A_{ij} x^i x^j + 2A_{in} x^i x^n + A_{nn} (x^n)^2 + 2A_i x^i + 2A_n x^n + 1 = 0.$$

Hyperquadrics cross the beginning of forming vector element.

In this case $A = 0$ and conditions of invariance of quadric (3.3.) have the form

$$(3.8) \quad \begin{aligned} \delta A_{\alpha\beta} - A_{\alpha\gamma} \pi_\beta^\gamma - A_{\beta\gamma} \pi_\alpha^\gamma &= \Theta A_{\alpha\beta}, \\ \delta A_\alpha - A_\gamma \pi_\alpha^\gamma &= \Theta A_\alpha. \end{aligned}$$

Describing (3.8.) obtain

$$(3.9) \quad \begin{aligned} \delta \Lambda_{ij} - \Lambda_{ik} \pi_j^k - \Lambda_{kj} \pi_i^k - \Lambda_{in} \pi_j^n - \Lambda_{nj} \pi_i^n &= \Theta \Lambda_{ij}, \\ \delta \Lambda_{in} - \Lambda_{kn} \pi_i^k - \Lambda_{nn} \pi_i^n &= \Theta \Lambda_{in}, \\ \delta A_{nn} &= \Theta A_{nn}, \\ \delta A_i - A_k \pi_i^k - A_n \pi_i^n &= \Theta A_i, \\ \delta A_n &= \Theta A_n. \end{aligned}$$

Considering the general form we put $A_n = 1$. Then $\Theta = 0$. Conditions of invariance of hyperquadric that crosses the beginning of vector element will be:

$$\delta A_{ij} - A_{ik} \pi_j^k - A_{kj} \pi_i^k - A_{in} \pi_j^n - A_{nj} \pi_i^n = 0,$$

$$(3.10) \quad \begin{aligned} \delta A_{in} - A_{kn} \pi_i^k - A_{nn} \pi_i^n &= 0, \\ \delta A_{nn} &= 0, \end{aligned}$$

$$\delta A_i - A_k \pi_i^k - \pi_i^n = 0.$$

The hyperquadric proper is taken by the equation:

$$(3.11) \quad A_{ij} x^i x^j + 2A_{in} x^i x^n + A_{nn} (x^n)^2 + 2A_i x^i + 2x^n = 0.$$

Some types of invariant hyperquadrics joint to vector field.

Invariant hyperquadric which is defined by the fundamental objects of the first and second order.

1. Analyze tensor $\Lambda_{n\beta}^\alpha$. In general case.

$$(4.1) \quad \Lambda = \det|\Lambda_{n\beta}^\alpha| \neq 0.$$

It permits to introduce values $v_{n\beta}^\alpha$, that their components are defined from the correspondence,

$$(4.2) \quad \Lambda_{n\beta}^\alpha v_{n\beta}^\alpha = \delta_\beta^\alpha.$$

Differential equations of tensor have the form

$$(4.3) \quad dv_{n\beta}^\alpha = v_{n\gamma}^\alpha \omega_\beta^\gamma - v_{n\beta}^\gamma \omega_\gamma^\alpha + v_{n\beta\gamma}^\alpha \omega^\gamma.$$

Build tensor

$$(4.4) \quad \Gamma_{\beta\gamma}^\alpha = v_{n\delta}^\alpha \Lambda_{n\beta\gamma}^\delta.$$

Its differential equations in case of fixation of main parameters obtain the form

$$(4.5) \quad \delta \Gamma_{\beta\gamma}^\alpha = \Gamma_{\beta\gamma}^\alpha \pi_\beta^\delta + \Gamma_{\beta\delta}^\alpha \pi_\gamma^\delta - \Gamma_{\beta\gamma}^\delta \pi_\delta^\alpha$$

due to tensor which is defined by the correspondence (4.4.) consequently build tensors

$$(4.6) \quad \begin{aligned} \Gamma_\alpha &= \Gamma_{\beta\alpha}^\beta, \\ \delta \Gamma_\alpha - \Gamma_\beta \pi_\alpha^\beta &= 0. \end{aligned}$$

$$(4.7) \quad \begin{aligned} \Gamma_{\alpha\beta} &= \Gamma_\gamma \Gamma_{\alpha\beta}^\gamma, \\ \delta \Gamma_{\alpha\beta} - \Gamma_{\beta\gamma} \pi_\alpha^\gamma - \Gamma_{\alpha\gamma} \pi_\beta^\gamma &= 0. \end{aligned}$$

If differential equations of tensors (4.6.) and (4.7.) have structure of equations (3.4.) the equations of invariant hyperquadric which is determined by the fundamental objects of the first and second order and it doesn't cross the beginning of definite vector element has the form

$$(4.8) \quad \Gamma_{\alpha\beta} x^\alpha x^\beta + 2\Gamma_\alpha x^\alpha + 1 = 0.$$

2. Invariant hyperquadric which is determined by the fundamental object of the second order.

Owing to tensor $\Lambda_{n\beta\gamma}^\alpha$ build tensors

$$(4.9) \quad \begin{aligned} \Lambda_{n\alpha\beta}^1 &= \Lambda_{n\alpha\beta}^\gamma \Lambda_{n\gamma}^1, \\ \delta \Lambda_{n\alpha\beta}^1 &= \Lambda_{n\gamma\beta}^1 \pi_\alpha^\gamma + \Lambda_{n\alpha\gamma}^1 \pi_\beta^\gamma. \end{aligned}$$

$$(4.10) \quad \begin{aligned} \Lambda_{n\alpha\chi}^2 &= \Lambda_{n\alpha\beta}^\gamma \Lambda_{n\gamma}^2, \\ \delta \Lambda_{n\alpha\beta}^2 &= \Lambda_{n\alpha\gamma}^2 \pi_\beta^\gamma + \Lambda_{n\gamma\beta}^2 \pi_\alpha^\gamma. \end{aligned}$$

If differential equations of tensors (4.9.) and (4.10.) have structure of equations (3.4.) equation of invariant hyperquadrics which is determined by the fundamental object of the second order and it doesn't cross the beginning of definite vector element and it has the form

$$(4.11) \quad \Lambda_{n\alpha\beta}^1 x^\alpha x^\beta + 2\Lambda_{n\alpha}^1 x^\alpha + 1 = 0.$$

$$(4.12) \quad \Lambda_{n\alpha\beta}^2 x^\alpha x^\beta + 2\Lambda_{n\alpha}^2 x^\alpha + 1 = 0.$$

Definition. Quadric (4.11) and (4.12) is called the main hyperquadrics.

Theorem 3. In case of transition from vector field of n-dimensional affine space A_n to vector field of n-dimensional space of affine connection the splitting of main hyperquadric takes place.

Conclusion. It has been constructed the splitting of the basic geometric images vector field (points, straights, hyperplanes and hyperquadrics) in transition from dimensional affine space to the space of affine connection. All investigations have been fulfilled in the moving coordinate system of zero order.

ЛІТЕРАТУРА

1. Амінов Ю.А. Геометрия векторного поля // М.: Наука, 1990. – 208 с.
2. Синцов Д.М. Работы по неголономной геометрии // Київ: Вища школа, 1972.
3. Бюпігес С.С. Геометрия векторного поля // Изв.АН СССР / Серия математика, 1948. Т. 10, №1.
4. Слухаев В.В. Эквиафинно-инвариантные неголономные поверхности потока жидкости // Труды Томского университета / Томск, 1967. – Т. 191, С.74–80.
5. Georghiev Gh. Observatii asupra geometriei afine diferentiale a cimpurilor de vectori // Academia republicii populare Romine / Lucralire constatuiriil ge-omertie topologie. – Lasi, 25 iunie, 1958, C.127–138.
6. Тадеєв П.О., Кравчук О.А. До геометрії векторного поля п-вимірного афінного простору // Вісник Київського університету / Серія фізико-математичні науки, 2006, В.4, С. 61–69.
7. Лаптев Г.Ф. Дифференциальная геометрия погруженных многообразий // Труды Московского математического общества / М., 1953. – Т.2, С. 275–382.
8. Норден А.П. Пространства аффинной связности // М.: Наука, 1975. – 432 с.

REFERENCES

1. AminovJ.A. Geometry of Vector Field // M. Science, 1990. – 208 p.
2. SintsovD.M. The Works on Non-holonomic geometry // Kyiv: Higher School, 1972.
3. BeushguesS.S. Geometry of Vector Field // Ed. of USSR / Serie: mathematics, 1948, V. 10, №1.
4. SluhayevV.V. Equiaffine invariant non-holonomic surfaces of the liquate stream // Works of Tomsk university / Tomsk, 1967. – V. 191, P.74-80.
5. Georghiev Gh. Observatii asupra geometriei afine diferentiale a cimpurilor de vectori // Academia republicii populare Romine / Lucralire constatuiriil ge-omertie topologie. – Lasi, 25 iunie, 1958, C.127–138.
6. TadeyevP.O., KravchukO.A. On Geometry of vector field of n-dimensional affine space // Journal of Kyiv University / Serie physical and mathematical sciences, 2006, V.4, P. 61–69.
7. LaptevG.F. Differential Geometry of manifolds // The Works of Moscow mathematical society / M., 1953. – V.2, P.275–382.
8. Norden A.P. Spaces of Affine Connection // M.: Science,1975. – 432 p.

Расщепление свойств геометрических образов, ассоциируемых с аффинным векторным полем

П. О. Тадеєв, О. А. Кравчук

Аннотация. Авторами построено расщепление основных геометрических образов векторного поля (точек, прямых, гиперплоскостей и гиперквадрик) при переходе от n-мерного аффинного пространства в пространство аффинной связности. Все исследования были выполнены в подвижной системе координат нулевого порядка.

Ключевые слова: векторное поле, расщепления геометрических образов, аффинное пространство, гиперплоскость, гиперквадрика.